

Western Arctic Shelfbreak Eddies: Formation and transport

Michael A. Spall, Robert S. Pickart, Paula S. Fratantoni, Albert J. Plueddemann

Woods Hole Oceanographic Institution

Woods Hole, MA 02543

April 16 2007

submitted to: Journal of Physical Oceanography

Corresponding Author: mspall@whoi.edu

Abstract

The mean structure and time-dependent behavior of the shelfbreak jet along the southern Beaufort Sea, and its ability to transport properties into the basin interior via eddies, are explored using high resolution mooring data and an idealized numerical model. The analysis focuses on springtime, when weakly stratified winter-transformed Pacific water is being advected out of the Chukchi Sea. When winds are weak, the observed jet is bottom trapped with a low potential vorticity core, has maximum mean velocities of $O(25 \text{ cm/s})$, and an eastward transport of 0.42 Sv . Despite the absence of winds, the current is highly time dependent, with relative vorticity and twisting vorticity often important components of the Ertel potential vorticity. An idealized primitive equation model forced by dense, weakly stratified waters flowing off a shelf produces a mean mid-depth boundary current similar in structure to that observed at the mooring site. The model boundary current is also highly variable, and produces numerous strong, small, anticyclonic eddies that transport the shelf water into the basin interior. Analysis of the energy conversion terms in both the mooring data and the numerical model indicates that the eddies are formed via baroclinic instability of the boundary current. The structure of the eddies in the basin interior compares well with observations from drifting ice platforms. The results suggest that eddies shed from the shelfbreak jet contribute significantly to the offshore flux of heat, salt, and other properties, and are likely important for the ventilation of the halocline in the western Arctic Ocean. Interaction with an anticyclonic basin-scale circulation, meant to simulate the Beaufort Gyre, enhances the offshore transport of shelf water and results in a loss of mass transport from the shelfbreak jet.

1 Introduction

The cold halocline of the interior western Arctic Ocean provides a barrier between the warmer, deeper waters of Atlantic origin and the sea surface, which is ice covered for most or all of the year. If these Atlantic waters were able to penetrate to the surface, there would be enough warm water to melt the ice cover. This would of course have enormous implications for the entire ecosystem of the Arctic as well as for air-sea exchange and global climate. It is now generally accepted that the halocline waters originate from the shelf regions (Aagaard et al., 1981; Muench et al. 2000). However, fundamental aspects of this exchange are not

well understood. For example, it is not clear where this exchange takes place, when during the year it occurs, or by what physical mechanisms it is accomplished.

One process that may play a role in halocline ventilation is eddy generation from the boundaries of the basin, where the most energetic circulation exists (Aagaard and Carmack, 1994). The interior of the Canada Basin is populated with small eddies of $O(20\text{ km})$ diameter. Measurements from drifting ice platforms suggest that the eddies are overwhelmingly anticyclonic ($> 90\%$). Eddies appear to be concentrated in the halocline (50-150 m depth), although both deeper (Manley and Hunkins, 1985; D’Asaro, 1988a) and shallower (Pickart et al., 2005; Plueddemann and Krishfield, 2007; Timmermans et al. 2007) eddies have been observed.¹ Eddies may be either warm or cold relative to the surrounding water, although cold-core eddies appear to dominate (Manley and Hunkins 1985). Measurements taken from drifting ice platforms suggest 1-2 eddies are found every 100 km of drift (Hunkins, 1974; Manley and Hunkins, 1985; Plueddemann and Krishfield, 2007), and it has been estimated that the basin is filled with 100-200 of these features at any one time (Plueddemann and Krishfield, 2007).

Hydrographic data suggest that the Canada Basin eddies are formed from distinct water masses, perhaps related to the winter and summer water masses on the Chukchi shelf (Pickart et al., 2005). The heat and salt transported seaward by the most commonly observed eddies are believed to be fundamental to the ventilation of the interior Arctic halocline (e.g. Muench et al. 2000). These eddies also represent a source of nutrients and zooplankton to the central basin (Llinas et al., 2007), and are potentially important for the offshore flux of organic carbon (Mathis et al., 2007). That far fewer of the eddies are found in the eastern Arctic suggests a source in the western basin (Manley and Hunkins, 1985).

The idea that anticyclonic eddies observed in the interior Canada Basin originate from hydrodynamic instabilities of the Alaskan Coastal Current in summer was put forward by Hunkins (1974), Hart and Killworth (1976), and Manley and Hunkins (1985). Their results are suggestive only, however, because their linear stability analysis was limited to very idealized, one dimensional flow with vertical shear typical of the boundary region. They did not consider a realistic structure of the boundary current or bottom topography, seasonality, or nonlinearities, all of which may influence the stability of the current and offshore eddy

¹It should be noted that near-surface eddies are difficult to detect from the historical ice camp data, so there may be a bias in these measurements.

flux. Moreover, the eddies produced from this current would be surface-intensified and contain warm water in their cores, and thus be unable to explain the source of the subsurface, cold-core anticyclones observed so prevalently in the interior.

D’Asaro (1988b) proposed that anticyclonic vorticity generated by a frictional sublayer of the boundary current in Barrow Canyon might provide a means to produce the near zero potential vorticity sometimes found in the anticyclones, and that eddies would be shed where the topography turns sharply to the right at the mouth of the canyon. Observations do not show such strong shear near the boundary (Münchow and Carmack, 1997; Pickart et al., 2005), and it is more likely that the low potential vorticity of the eddies is due to the convective origin of their water masses in the Chukchi Sea.

A series of papers by Chao and Shaw has explored various means of generating anticyclonic Arctic eddies. Early papers showed that localized regions of buoyancy loss in the interior can give rise to patches of low potential vorticity water, which then adjust to form anticyclone/cyclone pairs (Chao and Shaw, 1996, 1998). Stress applied to the upper-layer cyclone (representing ice interaction) leads to its decay, resulting in a mid-depth anticyclone. However, these experiments require forcing on very small scales (deformation radius) and of unrealistic strength in order to produce reasonably strong eddies. More recently, Chao and Shaw have explored boundary current sources for the eddies (Chao and Shaw; 2002, 2003a, b). This series of papers provides a source of dense water on the shelf and adds various effects, such as a strong offshore baroclinic current (meant to represent the Beaufort gyre), a cross-shelf flux, a canyon, and an offshore undercurrent (meant to represent the Beaufort Undercurrent; Aagaard, 1984). Under various circumstances, small numbers of cold-core, anti-cyclonic eddies are formed in these simulations (e.g. eddies associated with sinking of dense water in Barrow Canyon, see Chao and Shaw, 2003b). However, these authors have imposed an unrealistically strong westward-flowing Beaufort Gyre along the boundary, instead of the eastward-flowing shelfbreak jet that is now known to exist.

Historical measurements have revealed that Pacific-origin waters flow along the southern edge of the Canada Basin (e.g. Mountain et al., 1976; Aagaard, 1984). Recent observations, including those from the western Arctic Shelf-Basin Interactions (SBI) program, have elucidated this circulation. For example, it is now known that, in the absence of strong winds, an eastward-flowing boundary current exists along the edge of both the Chukchi and Beaufort Seas (Nikolopoulos and Pickart, 2007; Mathis et al., 2007; Llinas et al., 2007, see Fig. 1). It

is a narrow (order 10-15 km) jet, trapped to the shelfbreak, with different seasonal configurations. It is comprised largely of water that entered the Arctic through Bering Strait and subsequently reached the shelfedge via Herald Canyon in the west, and Barrow Canyon in the east (Fig. 1). More is known about the Beaufort shelfbreak jet because of a high-resolution mooring array that was maintained across the current during SBI (see Nikolopoulos and Pickart, 2007). There are three overall seasonal states of the current. During summer it is predominantly surface-intensified and advects Alaskan Coastal Water, i.e. it is the eastward extension of the Alaskan Coastal Current (see also Mountain et al., 1976; Paquette and Bourke, 1974). During fall and winter the current becomes bottom-trapped, but often reverses (flows westward) under strong easterly winds (Nikolopoulos and Pickart, 2007; Llinas et al., 2007). Then in spring, as the winds weaken, the current advects the cold, dense winter water formed the previous winter in the Chukchi and Bering Seas (Weingartner et al., 1998; Muench et al., 1988). In this configuration the current is bottom-trapped, with the strongest eastward flow in the vicinity of the shelfbreak.

In each of the above configurations the current is characterized by strong mesoscale variability. Some of this is wind driven (especially in fall and winter due to frequent storms, Pickart et al., 2007), but some appears to stem from internal instabilities of the current. Using a collection of historical hydrographic sections, Pickart (2004) argued that the Beaufort shelfbreak jet should be baroclinically unstable. The observed pattern of temperature variance is maximum at the offshore edge of the boundary current at the depth of the upper halocline, consistent with baroclinic instability. Furthermore, synoptic sections from SBI have captured eddies being spawned from the current along the Chukchi shelfedge (Pickart et al., 2005; Mathis et al., 2007). These observations demonstrate that (1) eddies can be formed from the boundary current over the steep shelfbreak topography away from Herald and Barrow Canyons; (2) the spatial scales of the boundary current and eddies is $O(10\text{ km})$; and (3) there is a seasonal cycle associated with the type of eddy that is formed from the boundary current.

In this study we use data from the SBI mooring array deployed across the Beaufort shelfbreak jet, observations of interior eddies from drifting platforms, and an idealized numerical model of the shelfbreak jet to investigate the mechanisms of eddy formation in the western Arctic. We focus on the spring time period when the current is advecting the coldest Chukchi/Bering winter water, and consider the “free jet” case when there is no wind

forcing (in both the observations and in the model). The characteristics and dynamics of this configuration of the boundary current are presented first using the SBI data. This is followed by the model investigation which focuses on the dynamics of eddy formation and the associated flux of mass and properties into the interior. Aspects of the model boundary current, and the structure of the eddies that it forms, are compared with the mooring data and with the interior ice station data.

2 Free-jet configuration of the Beaufort shelfbreak current

2.1 Mooring data calibration and processing

From August 2002 to September 2004 a high-resolution mooring array was maintained across the Beaufort shelfbreak and slope as part of the SBI program. The array was situated near 152°W , approximately 150 km east of Barrow Canyon (Fig. 1). It consisted of conductivity/temperature/depth (CTD) profilers to measure the hydrography, and acoustic Doppler current profilers (ADCPs) and acoustic current meters to measure velocity. The moorings were spaced 3-5 km apart on the upper part of the slope that sampled the Pacific component of the boundary current (Fig. 2, locations BS2–BS6). This portion of the array provided four synoptic vertical sections per day of the hydrographic and velocity structure of the boundary current (each CTD section took approximately 45 minutes to occupy). The sections were objectively gridded using Laplacian-Spline interpolation onto a grid with 2 km spacing in the horizontal and 5 m spacing in the vertical. The velocity data were de-tided using the tidal constituents obtained from the raw hourly ADCP data,² and an alongstream direction was objectively defined using the first year of data (see Nikolopolous and Pickart, 2007). The positive alongstream direction is 125°T , which is generally aligned with the topography upstream (west) of the array. The positive cross-stream direction is oriented offshore at 35°T .

The profiling CTD sensors were calibrated using stationary CTD sensors located at the bases of the moorings, together with shipboard CTD casts taken during the deployment and recovery cruises. The stationary sensors were themselves calibrated during the recovery

²The dominant tidal signals were weak, less than 2 cm/s.

cruises by attaching the sensors to the shipboard CTD package and performing comparison casts in deep water. The calibration and processing procedures are described in Fratantoni et al. (2006). The accuracy of the sensors was estimated to be .002 C for temperature and better than .03 for salinity. Due to potential ridging of the pack-ice, the top-floats of the moorings were positioned to be at 40 m. Although the velocity profiles extended shallower than this, the vertical sections were limited to the part of the domain where the hydrographic and velocity measurements overlapped, hence the top 40 m are blank. This is not a problem for the present study since the boundary current is bottom-intensified during the time period of interest.

Sections were constructed of potential temperature (θ), potential density (σ_θ , referenced to the sea surface), salinity (S), alongstream (u) and cross-stream (v) velocity, and the different components of the Ertel potential vorticity,

$$\pi = \frac{-f}{\rho_0} \frac{\partial \sigma_\theta}{\partial z} + \frac{1}{\rho_0} \frac{\partial u}{\partial y} \frac{\partial \sigma_\theta}{\partial z} - \frac{1}{\rho_0} \frac{\partial u}{\partial z} \frac{\partial \sigma_\theta}{\partial y}, \quad (1)$$

where f is the (constant) Coriolis parameter ($1.45 \times 10^{-4} \text{ s}^{-1}$), and ρ_0 is the reference density ($1.028 \times 10^3 \text{ kg m}^{-3}$, calculated using the mean array data). The first term on the right hand side of (1) is the vertical stretching term, and the second term is the relative vorticity. Because we are working in a rotated coordinate system aligned with the general direction of the flow, the relative vorticity is dominated by the u_y term. The third term is the tilting vorticity (see Hall (1994) for a derivation of this term), which becomes important when the isopycnals slope strongly across the current (which happens in the Beaufort shelfbreak jet, see below).

2.2 Seasonal water masses

As mentioned in the introduction, there are three main seasonal configurations of the shelfbreak current. The velocity structure of each of these states is presented in Nikolopoulos and Pickart (2007). The seasonal timing of the water masses can be seen in Fig. 3, which shows the temperature distribution near the center of the current for the first year of data (site BS3 in Fig. 2). The warm Alaskan Coastal Water is clearly evident in late-summer/early-fall of 2002, and again in July 2003 (note that it arrived much earlier the second summer). Another

notable aspect of the figure are the deep, warm “spikes” that are prevalent in late-fall (but occur at various other times of the year as well). These are the signature of storm events when Atlantic Water is upwelled due to strong easterly winds (Münchow et al., 2006; Pickart et al., 2007). Notably absent from Fig. 3 is a clear signature of the Chukchi/Bering Summer Water, which is resident over the Chukchi Sea in the late summer months (Coachman et al., 1975). However, in the second year of data this water was present in significant amount in late-fall after the passage of the warm Alaskan Coastal Water (attesting to interannual variability of the hydrographic structure of the current).

The period that we are focusing on in this study is the time when the bulk of the coldest Chukchi/Bering winter water flowed by the array. In 2003 this occurred during the months of April–June (Fig. 3) when the temperature was $< -1.7^{\circ}\text{C}$, near the freezing point. Based on θ - S values and nutrient concentrations, this is the same water mass that often occupies the center of cold-core anticyclones observed in the southern Canada Basin (Muench et al., 2000; Pickart et al., 2005; Llinas et al., 2007). Using our vertical sections we constructed timeseries of the properties of the winter water in the boundary current during this period by integrating the area within the -1.7°C contour for each realization. The salinity of the winter water so computed shows considerable variation over the three-month period, ranging between 32 and 34 (Fig. 4a). This means that the boundary current is capable of ventilating both the lower and upper halocline. During the latter half of April the winter water was saltiest, with much of it corresponding to the “cold, hypersaline mode” that Weingartner et al. (1998) observed in the Chukchi Sea. This implies that dense polynya water from the northeast Chukchi shelf can end up in the shelfbreak boundary current. From about mid-May onward the winter water was significantly less dense, and, according to Melling’s (1998) definition, was lighter than the upper halocline of the Canadian Beaufort Sea (Fig. 4a). However, this density level is within the lower part of the “summer Pacific halocline” discussed by Steele et al. (2004).

The grey shading in Fig. 4a represents the range in salinity within the patch of water colder than -1.7°C for the collection of vertical sections. At times there is a wide range of winter water present (sometimes spanning the entire halocline, e.g. in mid-April). This is partly due to the amount (cross-sectional area) of winter water contained in the section, which is plotted versus time in Fig. 4b. Note that there are times when greater than 50% of the section (to 300 m) is occupied by the winter water, although after the first week of June the amount was generally less than 5%. The overall message from Fig. 4 is that, downstream

of Barrow Canyon, the boundary current advects the coldest winter water sometime after the conclusion of winter (this was true in the second year of data as well), and that there can be a significant range in the density and amount of winter water during a given year, spanning most of the halocline.

2.3 Structure and dynamics

It is our intent to investigate the variability of the boundary current, and its ability to form eddies, in the absence of external forcing. While the winds over the Chukchi and Beaufort Seas generally weaken in spring, and the frequency of storms decreases, there are still wind events that disrupt the current. Upwelling events tend to reverse the current, and downwelling events enhance it. One can see in Fig. 3 that there were two strong upwelling events during spring 2003 (one in April and one in May). In order to isolate the time periods when the boundary current was a free jet, we developed an objective procedure for identifying the wind events. Essentially, an upwelling (downwelling) event was identified when the following criteria were met: (1) significant easterly (westerly) winds measured at the Pt. Barrow weather station; (2) the dominant flow throughout the section was westward (strongly eastward); and (3) the average near-bottom salinity in the vicinity of the shelfbreak reached a value well above (below) the mean monthly value, due to the up-slope (down-slope) movement of the water. We used salinity rather than temperature because it increases monotonically with depth (during certain springtime storms cold water was upwelled). Note that our wind filter removes only the synoptic weather events, it is insensitive to the mean, large-scale winds that force the anticyclonic Beaufort Gyre circulation.

Using these criteria we identified 10 upwelling events over the time period April to June (not all of them strong) which were removed from our collection of sections (Fig. 4b shows these events).³ Since the hydrographic response to the wind tended to lag the current response, it meant that some velocity sections were removed even though the current was flowing eastward (shortly after the storm ended). In all, 161 realizations, or 44% of the data, were discarded over the three-month period (there were no downwelling events). This is unfortunate, since among other things it reduces the confidence of the computed statistics, but we are left with no choice in order to isolate the free jet. The well-behaved mean fields

³An additional current reversal was removed in May despite the presence of weak winds at the time.

and interpretable variability presented below gives us confidence that the remaining 203 realizations provide meaningful results.

Mean fields

The mean structure of the free jet is shown in Fig. 5. The presence of the cold winter water is evident at the shelfbreak (Fig. 5a), coincident with an eastward velocity maximum exceeding 20 cm/s (Fig. 5b). The mean temperature structure over this time period is quite different from the year-long mean section due to the absence of the Atlantic Water influence near the shelfbreak (since the upwelling storms have been removed). The velocity structure is different as well: the mean unforced jet is 75% stronger than the year-long average flow, and the free-jet does not have the deep tail that forms during the winter storm season (see Nikolopoulos and Pickart, 2007). The transport of the free jet over the three-month period is 0.42 Sv ($1 \text{ Sv} = 10^6 \text{ m}^3 \text{ s}^{-1}$). This is three times larger than the year-long average Pacific water transport of 0.14 Sv (Nikolopoulos and Pickart, 2007).

The potential vorticity structure of the boundary current provides important insights regarding the dynamics of the flow. For most large-scale ocean currents (and even for many energetic currents), the stretching vorticity is the dominant term in π . Because of the small spatial scales of the high-latitude Beaufort shelfbreak jet, however, one might expect the other two terms to play an important role in the space-time evolution of the current. Fig. 5c shows the ratio of the relative vorticity to the stretching vorticity for the mean free jet (same as $-\overline{u_y}/f$). One sees the enhanced cyclonic (anti-cyclonic) vorticity on the seaward (shoreward) side of the jet—although for the mean current the values are relatively small. The stretching vorticity (Fig. 5d) shows a pronounced minimum at the shelfbreak associated with the weakly stratified winter water, which was likely formed via convective overturning in the Chukchi Sea. Note the spreading of the 26.2 and 26.6 isopycnals, which bound the mean winter water, as they progress onshore (Fig. 5a). The thermal wind shear associated with this isopycnal spreading dictates the vertical extent of the current, and also gives rise to the two regions of enhanced tilting vorticity above and below the jet (Fig. 5e; note that, for a geostrophic flow, the tilting vorticity is a negative-definite quantity, displayed here as the ratio between the tilting and stretching terms). In the mean, the tilting term is also fairly small (comparable to the relative vorticity).

The average total Ertel potential vorticity is shown in Fig. 5f. For the mean jet it is qualitatively similar to the stretching vorticity. The distribution of π shows a very important feature of the Beaufort shelfbreak jet; namely, that the cross-stream gradient of potential vorticity changes sign with depth. At the depth level of the jet core π increases offshore, from the low values in the winter water to the higher values of the ambient interior water. Below the jet core, the halocline is compressed due to the presence of the winter water pycnostad (note the 27 and 27.2 isopycnals in Fig. 5a), which results in a region of high potential vorticity adjacent to the slope near 180 m. This compression lessens offshore of the winter water, resulting in a decrease of π at this depth level. Such a change in the sign of π_y is a necessary condition for baroclinic instability of the current, which, as seen below, appears to lead to eddy formation.

Temporal variability

Despite the well-behaved mean fields in Fig. 5, the unforced jet is highly variable in time. This is seen, for example, in Fig. 4b, which shows that the amount of winter water measured by the array changes considerably independent of the wind, on a variety of timescales. To demonstrate this we present a synoptic snapshot of the current in late-April, when there was a large amount of winter water present (Fig. 6). Comparing this to the mean state (Fig. 5), one sees that, instantaneously, the boundary current can have a significantly different structure. In this realization the newly-ventilated winter water (colder than -1.75°C) occupies a large part of the slope, and the isopycnals beneath the pycnostad descend downward very sharply (Fig. 6a). The associated thermal wind shear is strong enough to reverse the jet near the bottom (Fig. 6b). The eastward-flowing portion of the jet is > 50 cm/s, with a relative vorticity that exceeds $.4f$ (Fig. 6c). While the stretching vorticity (Fig. 6d) shows the same feature as the mean—i.e. a region of low vorticity associated with the uniform winter water (100-150 m)—it also contains low values farther offshore where the isopycnals descend (150-200 m). The isopycnal slope is in fact so steep here that the tilting vorticity exceeds the stretching vorticity (Fig. 6e). Combined with the negative relative vorticity associated with the deep reversal of the jet, this results in a region of negative Ertel potential vorticity (Fig. 6f, near $x=30\text{-}35$ km). It is clear, therefore, that all three terms of π can be important synoptically in the Beaufort shelfbreak current.

How often is this the case? To answer this we created vertical sections of the ratio R_{pv} , defined as the sum of the relative plus tilting vorticity divided by the stretching vorticity.

As mentioned earlier, for large-scale flows $|R_{pv}| \ll 1$, and it is small for the mean free-jet of Fig. 5. However, based on Fig. 6 we expect that instantaneously R_{pv} will be significant. For each realization we determined the positive and negative extrema of R_{pv} , and where these values occurred in the vertical plane. Note that negative values of R_{pv} can result from both the tilting vorticity and anti-cyclonic relative vorticity, whereas positive values are due only to cyclonic relative vorticity.

Fig. 7a shows a histogram of the extrema of R_{pv} for the three-month period. The positive values are sharply peaked, with only 3% of the realizations containing $R_{pv} > .5$. By contrast, the negative values are broadly distributed over a large range, and more than 70% of the realizations contained $R_{pv} < -.5$. As seen in the example above (Fig. 6), the deflection of isopycnals due to the winter water can be large enough to make the total potential vorticity negative, and Fig. 7a indicates that this happened (i.e. $R_{pv} < -1$) a third of the time during the spring of 2003. Fig. 7b shows where the extreme values of R_{pv} occurred in relation to the mean jet. Consistent with the mean section (Fig. 5), the largest positive values occurred on the seaward side of the jet. There is, however, a second cluster of large positive values well offshore (near $x=33$ km). These correspond to periods when at least a portion of the jet was detached, or eddies were passing by the array. The negative values of R_{pv} occur in two general clusters, associated with the upward and downward deflection of isopycnals surrounding the winter water pycnostad (this is also consistent with the mean section). Interestingly, the largest negative values occur below the jet—which has strong ramifications for the eddy buoyancy flux and energetics of the current, as discussed later in the paper.

3 An idealized model of eddy formation and ventilation

The observations presented above have motivated us to consider the ability of a mid-depth boundary current of low potential vorticity water (the Chukchi/Bering winter water) to ventilate the western Arctic halocline through the formation of anticyclonic eddies. The presence of these eddies near the boundary current, in regions away from the two canyon outflows, suggests that they are formed all along the boundary, and that the topography associated with the canyons may be of second order importance. Our objectives in developing the model are to explore the efficiency of eddy formation from such a boundary current, to

compare the mean structure and variability of the boundary current and eddies to observations, and to assess the ability of this eddy formation mechanism to play an important role in the ventilation of the western Arctic halocline. Our approach is very idealized, allowing for a determination of the sensitivity of the eddy formations to both physical and numerical parameters. However, the central calculations are carried out in a parameter space representative of the observed circulation along the shelfbreak of the Chukchi and Beaufort Seas.

3.1 Model configuration

The numerical model used is the MIT general circulation model (Marshall et al. 1997). This model solves the hydrostatic, primitive equations on a uniform Cartesian, staggered C-grid with level vertical coordinates. A partial cell treatment of the bottom topography is accurate for steep topography in the presence of stratification, expected to be important for the present problem.

The model domain consists of a narrow island with sloping topography extending into deep water, and a shelf region 70 m deep at the western end of the island⁴ (See Fig. 8). The topography slopes linearly from 70 m depth to the maximum model depth of 450 m over a horizontal length scale of 19 km, giving a slope of $s = 0.02$ (this will be varied in Section 4.3). Areas of land are indicated by black. The horizontal resolution is 1 km and there are 27 levels in the vertical with spacing 10 m between 0 and 220 m depth and ranging from 30 m to 50 m between 220 m and 450 m. Since we are interested in the boundary current structure and eddies that exist within the halocline, the model domain is limited to the upper ocean. The initial stratification for temperature and salinity are uniform and taken from the May monthly mean PHC3.0 climatology (updated from Steele et al. 2001) at 150° W, 74° N.

The model incorporates second order vertical viscosity and diffusivity with coefficients $10^{-5} \text{ m}^2 \text{ s}^{-1}$. The vertical diffusion is increased to $1000 \text{ m}^2 \text{ s}^{-1}$ for statically unstable conditions in order to represent vertical convection. Horizontal viscosity is parameterized as a second order operator with the coefficient A_h determined by a Smagorinsky (1963) closure as

⁴Although all calculations here are on an f-plane, we refer to the direction increasing towards deep water as north and the direction along the island towards the right as east.

$$A_h = (\nu_s/\pi)^2 L^2 D. \quad (2)$$

ν_s is a nondimensional coefficient taken to be 1 here, L is the grid spacing, and D is the deformation rate, defined as

$$D = \left[(u_x - v_y)^2 + (u_y + v_x)^2 \right]^{1/2} \quad (3)$$

where u and v are the resolved horizontal velocities and subscripts indicate partial differentiation. For a deformation rate is typically dominated by the lateral shear, which can vary from $O(10^{-5} \text{ s}^{-1})$ for eddies in the interior to $O(10^{-4} \text{ s}^{-1})$ for the no-slip shear layer between the shelfbreak jet and the boundary. For the grid spacing of 1 km, this gives a range of lateral viscosity between $1 - 10 \text{ m}^2 \text{ s}^{-1}$.

The model is forced by generating dense water over a 100 km portion of the shelf in the western part of the domain (region of grey shading in Fig. 8) and advecting this water towards the shelfbreak. The water mass properties (temperature and salinity) and a barotropic northward flow are forced through the addition of restoring terms to the temperature, salinity, and meridional momentum equations. The restoring time scale is one hour and is applied only within the gray shaded region. Salinity is restored towards 30.5, 31.25, 32.0, 32.5, 32.5, 32.5, 32.5 over the 7 model levels over the shelf. Temperature is restored towards -1.8° C , and the meridional velocity towards 7.5 cm s^{-1} for the 70 m shelf depth. In addition, the model temperature and salinity are restored towards the initial climatological stratification within 50 km of the eastern and northern boundaries of the domain with a time scale of 5 days. This suppresses baroclinic Kelvin waves that are generated where the dense waters interact with the solid boundary at the eastern and western ends of the island and prohibits these waves from returning to the shelf region and contaminating the intended outflow. The model calculates density from a linear equation of state as

$$\rho = \rho_0 + \beta(S - S_0) - \alpha T \quad (4)$$

where $\beta = 0.8 \text{ Kg m}^{-3}$ and $\alpha = 0.04 \text{ Kg m}^{-3} \text{ C}^{-1}$ are the haline and thermal expansion coefficients and $\rho_0 = 1026.5 \text{ Kg m}^{-3}$ is a reference density and $S_0 = 32.5$ a reference salinity.

The model formulation is intended to represent the eastward flow of Pacific-origin winter water along the shelfbreak of the Chukchi and Beaufort Seas. The data described in Section 2 and in Pickart et al. (2005) suggest that eddies are formed all along the shelfbreak and are not confined only to the areas in the vicinity of Herald and Barrow Canyons. Although some previous modeling studies have considered the flow in the vicinity of such canyons (e.g. Chao and Shaw 2003b; D’Asaro 1988a), the focus of this study is on the stability of the boundary current away from these relatively small scale topographic features.

There are several advantages to this formulation. The northward transport is provided by water flowing anticyclonically around the island, and thus avoids the need for open boundary conditions. In addition, the structure of the boundary current is not specified but instead develops naturally as part of the solution. The stability of the boundary current is very sensitive to its potential vorticity structure and we wanted to avoid specifying this as an initial condition since this would directly control the outcome of the simulation. This approach also provides a continual source of kinetic and potential energy for the boundary current and thus avoids the spin-down that is inherent in periodic domain calculations of unstable currents. This idealized approach also allows us to explore the sensitivity of the eddy formation process to environmental parameters such as the bottom slope and physical processes such as ice-induced drag and lateral friction.

The Arctic Ocean is ice covered for much of the year near its margins, and year-round in the interior. It has been speculated that drag induced by the ice cover is important for the long-term evolution of eddies in the Arctic interior (Chao and Shaw, 1996; Ou 1986). In order to represent this stress, a quadratic drag law has been applied to the uppermost model level with coefficient $C_D = 2 \times 10^{-4}$. This gives a spin-down time scale of $\tau = H/C_D U^2$ which, for a thickness of $H = 50$ m and a horizontal velocity of $U = 10$ cm s⁻¹, gives $\tau \approx 25$ days. This will dampen near-surface velocities on the order of the boundary current strength fairly quickly compared to the estimated life time of the eddies in the absence of ice (note that this damping will have little impact on the anticyclonic eddies centered near 100 m depth).

The model is run for a period of 180 days. It takes about 40 days for the outflow from the model shelf to reach the eastern end of the island. The analysis in this section will focus on the mean and eddy variability of this central calculation, while the dynamical processes

involved in the eddy formation and its sensitivity to model configuration and parameters will be discussed in the following section.

3.2 Mean boundary current

The vertical structure of the mean boundary current is shown in Fig. 9, averaged over the final 100 days of integration and between 100 km and 700 km longitude. The zonal flow is sub-surface intensified with a maximum mean velocity of $O(25 \text{ cm s}^{-1})$. The maximum velocity is located several kilometers offshore of the sloping bottom, reflecting the no-slip lateral boundary conditions in the model. The net eastward transport is 0.37 Sv, similar to the mean transport of 0.42 Sv calculated from the mooring array during periods of weak winds.

The salinity within the eastward flowing boundary current is nearly uniform with a value close to 32.5, the same as the waters advected off the shelf at the western end of the island. The shallow isohalines slope upward towards the boundary, consistent with the mooring data presented above and with climatological observations (Pickart 2004). The deep isohalines slope downward towards the boundary, the spreading resulting from the low potential vorticity in the core of the shelfbreak jet. Thermal wind balances the eastward flowing, mid-depth maximum boundary current. The Ertel potential vorticity (1) is shown in Fig. 9b. Because of the very weak stratification on the shelf where the boundary current waters originated, the boundary current water has potential vorticity close to zero. There is a band of high potential vorticity in the upper 50 m over the jet, which results from the outcropping isopycnals and enhanced stratification. There is also a region of enhanced potential vorticity near 200 m depth, where the downward sloping isopycnals intersect the bottom. This general pattern is consistent with the mooring data (Fig. 5). The lateral gradient of potential vorticity is positive at mid-depth and negative in both the upper and lower boundary current, consistent with the mooring data and suggestive of baroclinic instability. There is a weak minimum of potential vorticity in the basin interior at approximately 120 m depth, close to the depth of the low potential vorticity boundary current water. This minimum is found in the PHC climatology, and is perhaps reflective of the transport of weakly stratified shelf-origin waters offshore.

3.3 Ventilation by anticyclonic eddies

As time progresses in the model simulation, the boundary current becomes unstable and sheds numerous mid-depth, anticyclonic, low potential vorticity eddies into the basin interior. After 24 days, the low potential vorticity waters have penetrated 500-600 km eastwards along the sloping bottom (See Fig. 10a). A series of meanders has developed and, in some cases, narrow filaments of low π water are extending into the interior. The relative vorticity within these filaments is anticyclonic and the relative vorticity between the filaments is cyclonic. The initial meander development is characterized by a dipole pair, or heton (Hogg and Stommel 1985), with a mid-depth anticyclone and a shallow cyclone slightly offset in the upstream direction. The cyclones are composed of ambient waters from the interior, and thus do not contain an anomalous potential vorticity or tracer signal. They are most evident in the pressure and velocity field, although filaments of low π boundary current water often get wrapped around their perimeters during the formation process making them identifiable in the tracer field as well. This configuration is typical of what is found for unstable currents (Pedlosky 1985; Spall, 1995; Bush et al. 1996; Shimada and Kubokawa, 1997) and is very effective at transporting tracers away from the front. After 180 days (Fig. 10b), the model has formed numerous low π , anticyclonic eddies that have propagated well into the basin interior. They are formed all along the boundary, beginning just downstream from the shelf region where the model is forced. The eddies vary in size but are typically 30 km in diameter with cores of nearly uniform, very low potential vorticity. The upper ocean cyclonic partners that had originally formed along with these mid-depth anticyclones have decayed as a result of the quadratic drag applied to the surface layer, parameterizing stress induced by interactions with the ice (consistent with the results of Chao and Shaw, 1996).

A vertical section on day 180 through the low π eddy located near 400 km longitude and 160 km latitude is shown in Fig. 11. This eddy was formed around day 60, propagated as far north as latitude 300 km, and then moved back towards the shelf. The velocity is anticyclonic throughout the upper 300 m with maximum velocities of $O(20 \text{ cm s}^{-1})$. The salinity shows the weak stratification in the core of the eddy, with a doming of isohalines above and a deepening of isohalines below the core of the eddy. The potential vorticity field reflects this low π core. Note that the anomalous potential vorticity signal is confined to the density range of the anomalous source waters, located between 90 m and 150 m. The velocity signal extends over a larger range of the water column because the waters both above and

below the low π core of the eddy are compressed as the weakly stratified core of the eddy passes by. This compression of the water column forces an anticyclonic circulation in order to conserve potential vorticity (Spall, 1995).

Fig. 11c shows tangential velocity vs. radius at a depth of 115 m for the eddy shown in the vertical slice in Fig. 11a,b. The tangential velocities at each model grid point are indicated by the black symbols, and the white line is a cubic spline fit to the model data points. The velocity has been scaled by the maximum of the spline fit profile ($V_{max} = 19 \text{ cm s}^{-1}$) and radius has been scaled by the radius of this maximum tangential velocity ($R_{max} = 12 \text{ km}$ where $V = V_{max}$). The degree to which the eddy is radially symmetric is evident in the scatter of the individual model grid points. The eddy core ($r < R_{max}$) shows relatively less scatter and is consistent with solid-body rotation. The region of velocity decay ($r > R_{max}$) shows increasing variability with increasing distance from the eddy center.

The velocity magnitude and radial structure of the model eddy are consistent with that observed from drifting platforms. Direct current observations reported by Manley and Hunkins (1985) indicate values of V_{max} from 30 - 50 cm/s. D’Asaro (1988b) reports V_{max} from 10 - 30 cm/s for three different halocline anticyclones, Plueddemann and Krishfield (2007) report statistics from 29 eddy encounters where the mean V_{max} was 24 cm/s. Thus, the strength of the model eddy ($V_{max}=19 \text{ cm/s}$) is representative of, although perhaps slightly weaker than, the typical observed eddy. The model produces eddies that range in radius from approximately 5 km to 20 km, but all show a similar velocity profile when scaled the same way.

Despite many Arctic eddy observations, reports of tangential velocity vs. eddy radius, passing from near the eddy center through the radius of maximum velocity, are relatively rare. Where radial structure has been reported (Newton et al., 1974, D’Asaro, 1988b, Timmermans et al., 2007) it is consistent with a solid-body core and a decay region of about 1.5 times R_{max} . Observations from ADCPs on ice-tethered drifters (Plueddemann and Krishfield, 2007) were used to produce a normalized radial structure to compare with the model eddy. Velocity data from 29 eddy encounters that clearly passed into the solid-body core were rotated into radial and tangential components using an estimated eddy center, and a third order polynomial fit was made assuming $V = 0$ at $R = 0$. Tangential velocity and radius from the fits were then scaled in the same manner as described above for the model eddy. The average of the normalized radial structure from the polynomial fits is shown in Fig. 11c, and

compares very well with the radial average structure of the model eddy. The observations indicate a self-similar structure for the 29 eddies, typical values in strength of 20 to 35 cm/s and in size from 2 to 9 km. Although the model eddies are somewhat larger and weaker, the good agreement in normalized radial structure with the observations is encouraging.

The anticyclonic eddies in the model transport waters that originated on the shelf into the basin interior. This is evident from the salinity and potential vorticity signals in the core of the eddies. A passive tracer has been added to the calculation in order to quantify the offshore transport by the eddies. This tracer is advected and diffused in the same way as are temperature and salinity, but is initialized at zero throughout the domain and maintained at a value of 1 within the forcing region on the shelf. A plan view of the tracer on day 180 at depth 115 m is shown in Fig. 12. The tracer clearly marks the location of the boundary current and core of the offshore eddies. The offshore flux of boundary current water, normalized by the flux off the shelf, is quantified by integrating the tracer flux as

$$F = \frac{\int_0^X \int_0^H v(y_0) T_p(y_0) dz dx}{\int_0^{X_s} \int_0^H v(y_s) T_p(y_s) dz dx} \quad (5)$$

where T_p is the passive tracer, H is the bottom depth, v is the meridional velocity, $X = 700$ km is the zonal extent of the integral for the offshore flux, $X_s = 100$ km is the western end of the shelf region, $y_s = 60$ km is the northern limit of the island, and $y_0 = 110$ km is the latitude where the offshore flux is calculated. A similar calculation is applied along a meridional section at $x = X$ between $y = y_s$ and $y = y_0$ to calculate the amount of tracer that is carried out of the region in the boundary current. The offshore and eastern locations of the flux calculation are indicated on Fig. 12 by the white lines. The offshore and alongshore fluxes are shown in Fig. 13 as a function of time. The offshore flux averaged between day 80 and day 180 is 53% of the flux off the shelf (indicated by the bold line). The flux is always offshore, but is characterized by strong, high frequency events resulting from individual eddy formations. The average of the sum of the alongshore flux and the offshore flux accounts for 93% of the flux off the shelf over this same time period, so there is little temporal storage within the integration contour on these time scales, although there can be significant storage on synoptic time scales. The offshore and alongshore fluxes are also generally out of phase, indicating that the offshore flux occurs at the expense of the

alongshore flux, even on synoptic time scales. Thus, the eddies are capable of transporting a significant amount of the shelf water into the basin interior.

4 Discussion

The standard model configuration presented above develops a mid-depth, low potential vorticity boundary current that readily forms anticyclonic eddies which ventilate the interior with shelf water. The mean structure of the model boundary current compares well with the free-jet configuration of the Beaufort shelfbreak jet obtained from the mooring data, and the eddies that it spawns have similar structure to those observed in the southern Canada Basin from drifting platforms. In this section, the mechanisms of the boundary current instability, its impact on the large-scale circulation and boundary current transports, and its sensitivity to model parameters are explored.

4.1 energetics

Model boundary current

The source of the boundary current instability is indicated by the terms that convert mean energy into eddy energy. The conversion from mean potential energy into eddy energy is often related to baroclinic instability and is given by

$$BC = -g\gamma\overline{v'\sigma_\theta'} / \rho_0 \quad (6)$$

where an overbar indicates the time mean, the primes indicate deviations from the time mean, and $\gamma = \partial z / \partial y$ is the mean isopycnal slope, and ρ_0 is a reference density. Potential energy is extracted from the mean flow by eddies transporting density down the mean density gradient. The cross-stream eddy density flux $\overline{v'\sigma_\theta'}$ calculated over the final 100 days of integration and then averaged between 100 km and 700 km is shown in Fig. 14a, superimposed on the mean density field. There is a positive density flux in the upper 100 m beginning on the offshore side of the mean boundary current and extending into the basin interior. Beneath this, deeper than 100 m, is a similar negative eddy density flux. This is the signal of eddies transporting water of shelf origin offshore into the stratified interior. The weak decay of the

eddy density flux offshore of the maximum is a result of the eddies remaining as coherent features that transport temperature and salinity into the interior.

The BC energy conversion calculated using this eddy density flux and the mean density field is shown in Fig. 14b. A positive value indicates that energy is being extracted from the mean by the growing eddies. The primary regions of energy conversion are centered on the sloping isopycnals near 50 m and 150 m depth on the offshore side of the boundary current, i.e. where the isopycnals spread due to the shelf water pycnostad on the slope (Fig. 9b). It is important to note that the energy conversion is limited to the region of sloping isopycnals even though the eddy density flux continues far offshore.

The conversion from mean kinetic energy into eddy energy is related to barotropic instability and is given by

$$BT = \overline{v'u'}U_y \quad (7)$$

where U is the time mean zonal velocity. Kinetic energy is extracted from the mean flow by eddies transporting zonal momentum down the mean velocity gradient. The eddy momentum flux $\overline{v'u'}$ calculated over the final 100 days and then averaged between 100 km and 700 km is shown in Fig. 14c. The maximum signal is centered near 100 m depth and extends approximately 20 km offshore. The sense of the eddy momentum flux is such that zonal momentum is transported offshore, leading to an offshore shift in the location of the boundary current. This is consistent with the offshore shift of the depth integrated transport streamfunction as the flow moves eastward along the island (discussed below). The change in zonal momentum that would result from this magnitude of eddy momentum flux divergence, over the zonal extent of the island with mean boundary current strength of 10 cm s^{-1} , is $O(10 \text{ cm s}^{-1})$. This would cause the mean flow onshore of the maximum eddy momentum flux to decrease by this amount and the mean flow offshore of the location of the maximum to increase by this amount, again consistent with the offshore shift found in the model.

Unlike the eddy density flux, which decays only weakly offshore of its maximum (Fig. 14a), the eddy momentum flux decreases rapidly in the meridional direction, resulting in weakly negative momentum flux offshore of the mean boundary current location (Fig. 14c). This means that while salinity (and other tracers) are carried far from the boundary by the eddies, momentum is not. Hence, the volume transport of the boundary current remains essentially

trapped near the boundary, even as the tracers originally within the current are carried far offshore. Due to the symmetry of the eddies, it is clear that they can not transport momentum but, due to the antisymmetry in density of the baroclinic eddy pairs, they are very effective at transporting density. The eddies are only able to transport momentum during their growth phase, not after they have separated from the mean current, which explains why the current remains close to the boundary.

The energy conversion from the mean kinetic energy to eddy energy is shown in Fig. 14d. The current is also barotropically unstable with the main region of energy conversion near the core of the boundary current at the depth of weakest stratification. There is also a region of weak energy conversion near the surface. The sense of the mid-depth conversion is to extract energy from the mean, although the integrated conversion over the entire section is about an order of magnitude less than the integrated baroclinic conversion term. Therefore, the instability of the model boundary current is most aptly described as a baroclinic instability.

Measured boundary current

The eddy fluxes of density and momentum, and the baroclinic and barotropic conversion terms, were also calculated from the mooring data. The mooring array does not extend as far offshore or as shallow in the water column, but the measurements do cover the core of the jet and allow for some comparisons with the energetics in the model. (The approximate region where the data and model overlap is indicated by the white lines in Fig. 14.) As in the model, the observed eddy density flux is positive in the upper portion of the shelfbreak jet and negative in the lower portion, although the zero-line is tilted upwards towards the boundary in the data whereas the transition from positive to negative is at the mid-depth of the model jet (Fig. 15a). The magnitude of the flux calculated from the data is similar to that produced in the model. The observed baroclinic conversion term is strongly positive beneath the jet where the isopycnals slope downward (near 150 m, Fig. 15b). This is consistent with the model also and indicates conversion from mean potential energy into eddy energy. There is, however, a second area of strong baroclinic conversion in the model above the jet where the isopycnals slope upward (near 60 m depth), which is difficult to resolve conclusively in the observations (although the measured conversion does become positive again at these depths, and data shallower than 40 m are not available).

The observed eddy momentum flux is mostly positive in the upper 150 m with the maximum located at the offshore limit of the array, approximately 20 km from the shelfbreak. The magnitude is similar to that found in the model, although the maximum in the model is located closer to the topography. The data also show a strong negative eddy momentum flux below the mean jet, centered at 230 m in a region of weak mean horizontal velocity shear. While the momentum flux in the model is negative at this depth as well, the value is close to zero. Both the data and model have positive barotropic conversion of similar magnitude near the depth of the jet (details of the distribution vary), indicating energy transfer from the mean kinetic energy to eddy energy. Integrating over the entire section, the observed baroclinic conversion is an order of magnitude larger than the barotropic conversion, which was the case in the model as well. Overall then, despite some detailed differences between the model and data, the general patterns and strengths of the eddy fluxes and energy conversion are comparable. This strongly supports the conclusion that the anticyclones observed in the interior of the southern Canada Basin originate from the shelfbreak jet, and that they are formed primarily by baroclinic instability all along the boundary.

4.2 mean transport streamfunction

The mean depth integrated transport streamfunction over the final 100 days of integration is shown in Fig. 16a. The circulation is dominated by an eastward flow along the north side of the island and a cyclonic circulation over much of the interior. The circulation around the island is initiated by the northward flow off the shelf between 0 and 100 km longitude. As this dense water flows over deeper topography, the influence of rotation deflects the current towards the right, along the sloping bottom. As mentioned earlier, the total transport off the shelf is approximately 0.37 Sv. The off-shelf transport is indicated in Fig. 16a by the black contours originating on the shelf and turning sharply to the right west of 100 km longitude. This flow remains close to the boundary, although it moves towards deeper water and broadens slightly as it flows eastward along the sloping bottom. This is consistent with the divergence of the eddy momentum flux discussed above. The circulation just offshore of the boundary current is primarily cyclonic and of basin scale in the zonal direction. The circulation farther in the interior is both cyclonic (most of the domain) and anticyclonic (around the isolated eddy near 100 km longitude and 150 km longitude).

The year-long eastward transport of the Beaufort shelfbreak jet at the array site (Fig. 1) is 0.14 Sv (Nikolopoulos and Pickart, 2007). This is less than 20% of the long-term transport of Pacific water through Bering Strait (0.8 Sv, Roach et al, 1995), and suggests that most of the Pacific water entering the Arctic does not end up in the boundary current east of Barrow Canyon. Even considering periods of weak winds in spring, the focus of this paper, the along-shelf transport is still only $O(50\%)$ of the Bering Strait transport. As described above, our central model calculation shows a slight offshore shift of the eastward boundary current transport, but actually produces an enhanced eastward transport compared to what flows off the shelf. This would seem to be at odds with the mooring data. To investigate this, an additional calculation was carried out in which a wind stress with anticyclonic curl was applied to the basin interior, defined as

$$\tau^x = -\tau_0 \cos(\pi(y - y_s)/(y_n - y_s)) \quad (8)$$

where $\tau_0 = 0.005 \text{ N m}^{-2}$ and y_n is the northern latitude of the domain. This magnitude of wind stress was chosen to give a mean wind-driven circulation in the basin interior of $O(4 \text{ cm s}^{-1})$, consistent with the strength of the anticyclonic Beaufort Gyre (Plueddemann et al. 1998). The vorticity input by the wind is dissipated through quadratic bottom friction with coefficient $C_D = 10^{-3}$. This does not significantly influence the boundary current structure or eddy formations, but is necessary for balancing the vorticity input by the wind.

The mean transport streamfunction for this wind-driven calculation is shown in Fig. 16b. The interior circulation is now anticyclonic, as expected from the wind forcing. However, the transport near the sloping topography is now eastwards only over the shallow topography. As the boundary current water is shifted offshore due to the eddy momentum flux, it becomes entrained into the westward flowing wind-driven gyre. The gyre is sufficiently strong that it is able to reverse the direction of flow, and with it carry the boundary current water towards the western basin. This is evident by the black contours marking the shelf water transport in Fig. 16b. By approximately 600 km longitude, essentially all of the shelfbreak jet transport has been lost to the basin interior. Hence, the slight offshore diversion of the shelfbreak jet driven by the eddy fluxes, interacting with the interior wind-driven gyre, results in a major transport loss of the boundary current. The region of eastward, eddy-driven, flow in the southern interior (Fig. 16a) is also competing with the westward wind-driven flow so that

there is a broad region of only weak mean flow offshore of the boundary current, between approximately 50 km and 125 km latitude. The mean transport streamfunction found for this wind-driven case is very similar to what is found if the transport for the central calculation with no wind is simply added to the transport for a wind-driven flow with no dense water outflow.

The eddies form in the wind-driven calculation much like they did in the central calculation, although the eddies are advected towards the interior and west by the model Beaufort Gyre, as seen by the passive tracer field on day 180 in Fig. 17. This is consistent with the pathways of Bering Summer Water inferred from observations by Shimada et al. (2006). The eddy distribution away from the boundary current is significantly different for the wind-driven case (Fig. 17) compared to the standard case (Fig. 12). The wind-driven case is distinguished by a relative sparsity of isolated anticyclones in the eastern half of the domain and a concentration in the west, with many of the western eddies well offshore of the boundary current. The eddy distribution observed from drifting platforms is generally consistent with the wind-driven model distribution. In particular, drifters carried around the periphery of the Canada Basin by the Beaufort Gyre showed relatively few eddies to the north and east of the Alaskan Beaufort Shelf and more frequent eddy encounters directly offshore of the shelf and to the west of Barrow Canyon (Plueddemann et al., 1998; Krishfield et al., 2002). Although intriguing, this evidence is not definitive because the observations are only along the drift tracks and thus do not uniformly sample the region.

The average tracer flux carried into the interior over the final 100 days is increased to 80% of the flux off the shelf, significantly larger than the calculation without the wind. This is due, in part, to the advection by the wind-driven flow carrying the eddies away from the boundary in the western part of the domain. There is also now an offshore transport due to the mean flow because the boundary current transport is diverted offshore. The wind causes an enhanced horizontal shear of the velocity field due to the westward flow adjacent to the boundary current, however the energetics of this calculation look very much like that for the central calculation with no wind, so it does not appear that the instability process is strongly influenced by the wind.

4.3 parameter sensitivities

Lateral boundary condition and viscosity

The structure of the lateral shear of the boundary current is fundamental to the model boundary current stability and the formation of the eddies. A model calculation identical to the standard case except with free-slip lateral boundary conditions (instead of no-slip) produces no eddies or offshore flux of the passive tracer. The primary difference in the mean boundary current structure between these two calculations is the presence of a viscous shear layer near the boundary for the calculation with no-slip boundary conditions. The mean zonal velocity at 115 m depth, averaged over the final 100 days of integration and between 200 km and 700 km longitude, is shown in Fig. 18 for the free-slip and no-slip calculations. The velocity with free-slip increases towards the boundary across the eastward flowing boundary current to a maximum of approximately 60 cm s^{-1} with very little decay as the boundary is approached. The no-slip calculation has a much weaker maximum velocity, only approximately 20 cm s^{-1} , and decreases towards zero within about 2 km of the boundary (relative vorticity $\approx -0.7f$). The offshore relative vorticity is much larger for free-slip than it is for no slip, but it is positive throughout nearly all of the eastward flowing boundary current while it has a narrow region of large negative relative vorticity for no-slip boundary conditions. The isopycnal slope is of the same sign in both cases, although significantly steeper in the free-slip case. The change in stability is likely not a result of changes in the baroclinicity of the boundary current but instead related to the presence of the shear layer near the boundary. A similar destabilizing influence of lateral shear layers provided by no-slip boundary conditions for baroclinic instability has been found by Mundt et al. (1995) and Berloff and McWilliams (1999).

The size and growth rate of the eddies is also sensitive to the nondimensional viscosity parameter ν_s . Increased values result in larger eddies and an enhanced offshore transport of shelf tracer while smaller values result in smaller eddies and less offshore transport (see Table 1), although in all cases studied in the range of $0.75 < \nu_s < 1.5$ the model instability appears to be qualitatively similar. The shear layer on the on-shore side of the mean boundary current is the same width in all of these cases, $O(2 \text{ km})$, so the eddy scale is not simply related to the width of the shear layer.

Bottom slope

The percent of the passive tracer that gets carried into the basin interior by eddies is sensitive to the topographic slope between the shelfbreak and the interior. The offshore flux increases with increasing slope, largely due to the production of more eddies (Table 1). The offshore flux is less than 10% for a topographic slope of 0.01 and over 90% for a bottom slope of 0.03. This suggests that the offshore flux of boundary current water might vary strongly with location as the topographic slope changes along the boundary.

The topographic slope at the mooring array location is approximately 0.018, close to that used for the central model calculation. However, the main point of these calculations is to demonstrate which parameters of the model are most important for the formation of eddies and the offshore flux of shelf water. The important result is that realistic topographic slopes along the Chukchi and southern Beaufort shelfbreak fall within the range of topographic slopes for which we find significant eddy formations and offshore flux of tracer. However, because the offshore flux is also sensitive to the value of the viscosity coefficient ν_s , the effect of varying the bottom slope is relative; increasing the viscosity to $\nu_s = 1.5$ results in significant offshore tracer flux even when the bottom slope is relatively weak, $s = 0.01$ (Table 1).

Ice drag

The quadratic drag applied to the uppermost level in the model represents the interaction with ice cover, and damps the upper ocean motions with a time scale of $O(25 \text{ days})$. This affects most strongly the cyclonic eddy partners that are formed during the instability process and help to propagate the anticyclonic eddies away from the shelfbreak jet. Setting this drag to zero increases the offshore eddy flux by only 7%, while doubling it to 4×10^{-4} does not change the offshore flux at all (Table 1). Thus, the magnitude of the offshore flux is not very sensitive to small variations of the drag coefficient.

5 Summary

It has been known for some time that the southern Canada Basin is populated with a large number of cold-core, anti-cyclonic halocline eddies. While the properties of the eddies indicate a shelf source, the mechanism of formation has been unclear. We have demonstrated, using high-resolution mooring observations and an idealized numerical model, that such

eddies are likely spawned by the shelfbreak jet via baroclinic instability. Our analysis focused on the spring time period, when winter-transformed waters from the Bering and Chukchi Seas are carried eastward by the current. These waters are near the freezing point and very weakly stratified, forming a bottom trapped, low potential vorticity boundary current when the winds are weak. The salinity of these winter water masses ranges between 32–34, and are thus of interest for ventilation of both the upper and lower halocline. A mooring array 150 km east of Barrow Canyon measured an eastward transport of 0.42 Sv during periods when the boundary current is not strongly influenced by local winds. The maximum observed mean velocities were $O(25 \text{ cm s}^{-1})$, with a current width of $O(10 - 15 \text{ km})$. The current was highly time-dependent and nonlinear, with the relative vorticity and tilting vorticity terms often becoming important for the overall Ertel potential vorticity. The eddy density and momentum fluxes calculated from the array show significant cross-stream fluxes of both density and zonal momentum. The energy conversion terms associated with these eddy fluxes indicate that the current is both baroclinically and barotropically unstable, although the baroclinic conversion term is dominant.

An idealized numerical model forced by weakly stratified, dense water flowing off a shelf produces a boundary current in general agreement with the observations, both in its mean structure and in its time-dependent behavior. The model current is strongly unstable and produces numerous cold-core, anticyclonic, mid-depth eddies that transport the shelf-origin water into the basin interior. The eddy fluxes and energy conversion terms in the model compare reasonably well with those from the mooring data, although the mooring data do not cover the full offshore extent of the eddy formation region so a complete comparison is not possible. The structure of the eddies formed by the model compare well with typical velocity profiles of eddies measured in the basin interior by ADCPs from drifting ice stations. The model does show a sensitivity of the quantitative behavior (number of eddies formed, size of eddies) to details of the model configuration (bottom slope, frictional parameterizations), but the qualitative behavior is largely unchanged. The model is not intended to provide a detailed prediction of the shelfbreak jet system in the southern Beaufort Sea, but is instead used to demonstrate a plausible process and to identify parameters that are important for the stability of the shelfbreak jet. The results suggest regions in which large eddy-driven exchange can be expected (steep topography) and isolate physical processes that need to be better understood (shear layers and mixing near topography).

The positive comparisons between the model and the data, both in the region of eddy formation and in the basin interior, suggest that the model is faithfully representing the boundary current and its instability. The number of cold-core eddies estimated to exist at any one time in the basin interior, $O(100 - 200)$, together with their estimated lifetime of $O(1 - 2 \text{ years})$, indicates that approximately 100-200 eddies are formed each year (see also Pickart et al., 2005). Based on the 2-year mooring data set, winter water is present along the shelf break for roughly half of the year (keep in mind that we considered only the coldest winter water in this study), which requires the formation of an eddy approximately every 1-2 days. Such a large production rate seems unlikely to be supported by only one or two canyon outflows, but is not unreasonable based on the number of eddies formed in the model when the entire length of the shelf break is taken into consideration. The eddies in the central model calculation transport almost half of the original shelf water into the basin interior over a boundary extent of 600 km, indicating that the eddies are quite effective at transporting water masses and properties from the shelf region into the basin interior. This suggests that eddy formation is a major contributor to halocline ventilation in the western Arctic.

The mass transport of the boundary current, however, is much more closely confined to the boundary region than the tracer, because the eddies are not an effective means to transport momentum. However, when large-scale anticyclonic winds over the Beaufort Sea are included in the model, the slight offshore shift of the boundary current driven by the eddy fluxes is sufficient to get the boundary current transport entrained into the westward flowing wind-driven Beaufort Gyre. This results in a reduction of the eastward transport along the boundary and can result in a majority of the shelf water being transferred into the interior. One ramification of this is that an enhanced Beaufort Gyre might significantly redirect the Pacific-origin water into the interior of the Canada Basin. Steele et al. (2004) argue that the atmosphere-ocean conditions associated with the phase of the Arctic Oscillation (AO) strongly dictate the ability for Chukchi/Bering summer water to appear north of Ellesmere Island (and subsequently flow through Nares Strait and Fram Strait). They attribute this to a shift in the location and strength of the transpolar drift and Beaufort Gyre due to the AO. Our results suggest that the redirection of the boundary current may also play an important role (we note that Bering/Chukchi summer water is dense enough to flow eastward as a bottom-trapped boundary current). This is not inconsistent with Steele et al.'s scenario, but could contribute to it. Further work is required to flesh this out.

Acknowledgments

The authors wish to thank Anna Nikolopoulos and Daniel Torres for their help in processing and interpreting the mooring data. This study was supported by the National Science Foundation Office of Polar Programs under Grants 0421904 and 035268 (MS), and the Office of Naval Research Grant N00014-02-1-0317 (RP and PF). Analysis by AJP was supported by the Office of Naval Research under Grant N00014-97-1-0135 and by the National Science Foundation under Grant OPP-9815303. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation or the Office of Naval Research.

References

- Aagaard K. and Carmack E.C., 1994: The Arctic and climate. In: The Polar Oceans and Their Role in Shaping the Global Environment (edited by O.M. Johannessen, R.D. Muench and J.E. Overland, Geophysical Monograph Series, 85, AGU, Washington, D.C., pp. 4-20). AGU, Washington, D.C.
- Aagaard, K., L. K. Coachman, E. C. Carmack, 1981: On the halocline of the Arctic Ocean. *Deep-Sea Res.*, **28**, 529-545
- Aagaard, K., 1984: The Beaufort Undercurrent. *The Alaskan Beaufort Sea: Ecosystems and Environments*, Academic Press, pp. 47-71.
- Aagaard, K. and A. T. Roach, 1990: Arctic ocean-shelf exchange: measurements in Barrow Canyon. *J. Geophys. Res.*, **95**, 18,163-18,175.
- Berloff, P. S., and J. C. McWilliams, 1999: Quasigeostrophic dynamics of the western boundary current. *J. Phys. Oceanogr.*, **29**, 2607-2634.
- Bush, A. B. G., J. C. McWilliams, and W. R. Peltier, 1996: The formation of oceanic eddies in symmetric and asymmetric jets. Part II: Late time evolution and coherent vortex formation. *J. Phys. Oceanogr.*, **26**, 1825-1848.
- Chao, S-Y. and P-T Shaw, 1996: Initialization, asymmetry, and spindown of Arctic eddies. *J. Phys. Oceanogr.*, **26**, 2076-2092.
- Chao, S-Y. and P-T Shaw, 1998: Eddy maintenance and attrition in a vertically sheared current under Arctic ice. *J. Phys. Oceanogr.*, **28**, 2427-2443.
- Chao, S-Y and P-T Shaw, 2002: A numerical investigation of slanted convection and subsurface anticyclone generation in an Arctic baroclinic current system. *J. Geophys. Res.*, **107**, 10.1029/2001JC000786.
- Chao, S-Y and P-T Shaw, 2003a: A numerical study of dense water outflows and halocline anticyclones in an Arctic baroclinic current *J. Geophys. Res.*, **108**, 10.1029/2002JC001473.
- Chao, S-Y and P-T Shaw, 2003b: Eddy shedding from submarine-canyon plumes in an Arctic boundary current system: Sensitivity to the undercurrent. *J. Phys. Oceanogr.*, **33**, 2032-2044.

- Coachman, L.K., K. Aagaard, and R.B. Tripp, 1975: Bering Strait: The regional physical oceanography. *University of Washington Press*, Seattle, 172 pp.
- D’Asaro, E. A., 1988a: Observations of small eddies in the Beaufort Sea, *J. Geophys. Res.*, **93**, 6669-6684.
- D’Asaro, E. A., 1988b: Generation of submesoscale vortices: A new mechanism. *J. Geophys. Res.*, **93**, 6685-6693.
- Fratantoni, P.S., S. Zimmerman, R. S. Pickart, M. Swartz, 2006: Western Arctic Shelf-Basin Interactions Experiment: Processing and calibration of moored profiler data from the Beaufort shelf edge mooring array, Woods Hole Oceanographic Institution Technical report WHOI-2006-15, 34pp.
- Garrison, G. R and P. Becker, 1976: The Barrow submarine canyon: A drain for the Chukchi Sea. *J. Geophys. Res.*, **81**, 4445-4553.
- Hall, M.M., 1994: Synthesizing the Gulf Stream thermal structure from XBT data. *Journal of Physical Oceanography*, **24**, 2278–2287.
- Hart, J. E., and P. D. Killworth, 1974: Open ocean baroclinic instability in the Arctic. *Deep-Sea Res.*, **23**, 637-645.
- Hogg, N. G., and H. M. Stommel, 1985: The heton, an elementary interaction between discrete baroclinic geostrophic vortices and its implications concerning eddy heat-flow. *Proc. Roy. Soc. London*, **A397**, 1-20.
- Hunkins, K. L., 1974: Subsurface eddies in the Arctic Ocean. *Deep-Sea Res.*, **21**, 1017-1033.
- Llinas, L., R.S. Pickart, J.T. Mathis, and S.L. Smith, 2007: The effects of eddy transport on zooplankton biomass and community composition in the western Arctic. *Deep-Sea Research II*, submitted.
- Manley, T. O. and K. Hunkins, 1985: Mesoscale eddies of the Arctic Ocean. *J. Geophys. Res.*, **90**, 4911-4930.
- Marshall, J., C. Hill, L. Perelman, and A. Adcroft, 1997: Hydrostatic, quasi-hydrostatic, and non-hydrostatic ocean modeling. *J. Geophys. Res.*, **102**, 5733-5752.

- Mathis, J.T., R.S. Pickart, D.A. Hansell, D. Kadko, and N.R. Bates, 2007: Eddy transport of organic carbon and nutrients from the Chukchi shelf into the deep Arctic basin. *Journal of Geophysical Research*, in press.
- Melling, H., 1998: Hydrographic changes in the Canada basin of the Arctic ocean, 1979–1996. *Journal of Geophysical Research*, **103**, 7637–7645.
- Mountain, D. G., 1974: Bering Sea water on the north Alaskan shelf. Ph.D. Thesis, University of Washington, Seattle, WA, 153 pp.
- Mountain, D.G., L.K. Coachman, and K. Aagaard, 1976: On the flow through Barrow Canyon. *Journal of Physical Oceanography*, **6**, 461–470.
- Muench, R.D., J.D. Schumacher, and S.A. Salo, 1988: Winter currents and hydrographic conditions on the northern central Bering Sea shelf. *Journal of Geophysical Research*, **93**, 516–526.
- Muench, R. D., J. T. Gunn, T. E. Whitledge, P. Schlosser, and W. Smethie Jr., 2000: An Arctic Ocean cold core eddy. *J. Geophys. Res.*, **105**, 23997–24006.
- Münchow A., and E. Carmack, 1997: Synoptic flow and density observations near an Arctic shelfbreak. *J. Phys. Oceanogr.*, **27**, 1402–1419.
- Münchow, A., R.S. Pickart, T.J. Weingartner, R.A. Woodgate, and D. Kadko, 2006: Arctic boundary currents over the Chukchi and Beaufort slope seas: Observational snapshots, transports, scales, and spatial variability from ADCP surveys, *EOS Trans. AGU*, **87(36)**, Ocean Sci. Meet. Suppl., Abstract OS33N-03.
- Mundt, M. D., N. H. Brummell, and J. E. Hart, 1995: Linear and nonlinear baroclinic instability with rigid sidewalls. *J. Fluid Mech.*, **291**, 109–138.
- Newton, J. L., K. Aagaard, and L. K. Coachman, 1974: Baroclinic eddies in the Arctic Ocean. *Deep-Sea Res.*, **21**, 707–719.
- Nikolopoulos, A. and R.S. Pickart, 2007: The western Arctic boundary current at 152W: Structure, variability, and transport. *Deep-Sea Research II*, submitted.
- Ou, H. W. and A. L. Gordon, 1986: Spin-down of baroclinic eddies under sea ice. *J. Geophys. Res.*, **91**, 7623–7630.

- Paquette, R. G. and R. H. Bourke, 1974: Observations on the coastal current of Arctic Alaska. *J. Mar. Res.*, **32**, 195-207.
- Pedlosky, J. 1985: The instability of continuous heton clouds. *J. Atmos. Sci.*, **42**, 1477-1486.
- Pickart, R. S., 2004: Shelfbreak circulation in the Alaskan Beaufort Sea: Mean structure and variability. submitted to: *J. Geophys. Res.*.
- Pickart, R. S., T. J. Weingartner, L. J. Pratt, S. Zimmermann, and D. J. Torres, 2005: Flow of winter-transformed Pacific water into the Western Arctic. *Deep-Sea Res.*II, **52**, 3175-3198
- Pickart, R.S., G.W.K. Moore, A.M. Macdonald, J.E. Walsh, and W. E. Kessler, 2007: Seasonal evolution of Aleutian low-pressure systems: Implications for the North Pacific sub-polar circulation. submitted to: *J. Climate*.
- Plueddemann, A., R. Krishfield, and C. Edwards, 1999: Eddies in the Beaufort Gyre, Ocean-Atmosphere-Ice Interactions (OAI) All Hands Meeting, 20-22 October, 1999, Virginia Beach, VA.
- Plueddemann, A. J., R. Krishfield, T. Takizawa, K. Hatakeyama, and S. Honjo, 1998: Upper ocean velocities in the Beaufort Gyre. *Geophys. Res. Lett.*, **25**, 183-186.
- Plueddemann, A. J., R. Krishfield, 2007: Physical properties of eddies in the western Arctic. to be submitted to: *J. Geophys. Res.*
- Roach, A. T., A. Aagaard, C. H. Pease, S. A. Salo, T. Weingartner, V. Pavlov, and M. Kulakov, 1995: Direct measurements of transport and water properties through Bering Strait. *J. Geophys. Res.*, **100**, 18443-18457.
- Shimada, K. and A. Kubokawa, 1997: Nonlinear evolution of linearly unstable barotropic boundary currents. *J. Phys. Oceanogr.*, **27**, 1326-1348.
- Smagorinsky, J. 1963: General circulation experiments with the primitive equations: I. The basic experiment. *Mon. Wea. Rev.*, **91**, 99-164.
- Spall, M. A., 1995: Frontogenesis, subduction, and cross-front exchange at upper ocean fronts. *J. Geophys. Res.*, **100**, 2543-2557.
- Steele, M., R. Morley, and W. Ermold, 2001: PHC: A global ocean hydrography with a high quality Arctic Ocean. *J. Climate*, **14**, 2079-2087.

- Steele, M., J. Morrison, W. Ermold, I. Rigor, and M. Ortmeier, 2004: Circulation of summer Pacific halocline water in the Arctic Ocean. *J. Geophys. Res.*, **109**, doi:10.1029/2003JC002009.
- Timmermans, M.-L., J. Toole, A. Proshutinsky, R. Krishfield, and A. Plueddemann, 2007: Eddies in the Canada Basin, Arctic Ocean, observed from ice-tethered profilers. submitted to: *J. Phys. Oceanogr.*.
- Weingartner, T. J., D. J. Cavalieri, K. Aagaard, and Y. Sasaki, 1998: Circulation, dense water formation, and outflow on the northeast Chukchi shelf. *J. Geophys. Res.*, **103**, 7647-7661.
- Woodgate, R. A., K. Aagaard, and T. Weingartner, 2005: A year in the physical oceanography of the Chukchi Sea: Moored measurements from autumn 1990-1991. *Deep-Sea Res.*II, **52**, 3116-3149.

RUN	s	C_D	ν_s	F
1	0.02	2×10^{-4}	1.0	.53
2	0.02	2×10^{-4}	0.75	.11
3	0.02	2×10^{-4}	1.5	.91
4	0.01	2×10^{-4}	1.0	.07
5	0.03	2×10^{-4}	1.0	.93
6	0.02	4×10^{-4}	1.0	.53
7	0.02	0×10^{-4}	1.0	.60
8	0.01	2×10^{-4}	1.5	.73

Table 1: Normalized offshore tracer flux F for variations in Smagorinsky viscosity coefficient ν_s ; bottom slope s ; ice drag coefficient C_D .




Figure 1: Map of the region of interest and schematic circulation. A shelfbreak jet exists along the edge of the Chukchi and Beaufort Seas. In the Beaufort Sea it is fed by the Alaskan Coastal Current flowing through Barrow Canyon, and in the Chukchi Sea it is presumably fed by the outflow from Herald Canyon. Anti-cyclonic eddies are commonly observed offshore of the shelfbreak jet and in the Canada Basin within the Beaufort Gyre. The mooring locations of the SBI array are indicated.




Figure 2: Configuration of the SBI Beaufort slope profiling mooring array.




Figure 3: Timeseries of potential temperature ($^{\circ}\text{C}$) at the center of the boundary current (site BS3, see Fig. 2). The three month period of focus for this study is indicated.




Figure 4: Properties of the Chukchi/Bering winter water during spring 2003. The winter water is defined as that water colder than -1.7°C (see text). (a) Mean salinity (black line) and range of salinity (grey shading). The limits of the cold halocline based on historical data from the region (Pickart, 2004) are shown. (b) Area of the winter water expressed as a percent of the total area of the section to 300 m depth. The periods denoted by grey shading are upwelling events (see text).




Figure 5: Mean vertical sections of the unforced jet.




Figure 6: Vertical sections of a snapshot of the jet on 29 April, 2003 0600 GMT.




Figure 7: (a) Histogram of the positive and negative extrema of R_{pv} (see text) for the collection of sections. (b) Scatter plot of the extrema of R_{pv} overlaid on the mean alongstream velocity section of Fig. 5




Figure 8: Model topography (contour interval = 100 m) and domain. The black region is an island, the gray region is a 70 m deep shelf where the temperature, salinity, and momentum are forced.




Figure 9: Meridional section of the mean boundary current a) zonal velocity (cm/s); b) Ertel potential vorticity ($10^{-9} \text{ m}^{-1} \text{ s}^{-1}$). Isopycnals are indicated by the black contours.




Figure 10: Potential vorticity at 115 m depth on a) day 24 and b) day 180 ($10^{-9} \text{ m}^{-1}\text{s}^{-1}$). The white line in b) marks the location of the vertical section in Fig. 11




Figure 11: Vertical slice through the eddy indicated in Fig. 10b at the location of the white line on day 180. a) meridional velocity (colors, m/s) and salinity (contours); b) Ertel potential vorticity ($10^{-9} \text{ m}^{-1} \text{ s}^{-1}$); c) tangential velocity as a function of radius, nondimensionalized by radius of maximum velocity (12 km) and maximum velocity (19 cm s^{-1}). Circles mark each model grid point, the white line is a least squares fit through the model data points, and the red line is an average polynomial fit based on the observations of Plueddemann and Krishfield (2007).




Figure 12: Passive tracer in the model on day 180 at 115 m depth. The region of offshore and along shore tracer flux is indicated by the bold white lines. The tracer is initially zero everywhere in the domain, maintained at a value of 1 in the shelf forcing region, and is otherwise advected and diffused as temperature and salinity.




Figure 13: The offshore (solid line) and along shore (dashed line) advective flux of the passive tracer as a function of time, normalized by the flux of tracer off the shelf in the forcing region. The mean offshore tracer flux over the final 100 days of integration is indicated by the bold line (53%).




Figure 14: Along-shelf and temporal mean of a) offshore eddy density flux ($\text{Kg m}^{-2} \text{s}^{-1} \times 10^{-2}$); b) baroclinic energy conversion term ($\text{m}^2 \text{s}^{-3} \times 10^{-8}$); c) eddy momentum flux ($\text{m}^2 \text{s}^{-2} \times 10^{-3}$); d) barotropic energy conversion term ($\text{m}^2 \text{s}^{-3} \times 10^{-8}$), overlaid on the mean density field (thin black contours). The zero contour of each field is given by the bold black line. The white line indicates the region of overlap with the observations (Fig. 15).




Figure 15: Results from the mooring data for a) offshore eddy density flux ($\text{Kg m}^{-2} \text{s}^{-1} \times 10^{-2}$); b) baroclinic energy conversion term (6) ($\text{m}^2 \text{s}^{-3} \times 10^{-8}$); c) eddy momentum flux ($\text{m}^2 \text{s}^{-2} \times 10^{-3}$); d) barotropic energy conversion term (7) ($\text{m}^2 \text{s}^{-3} \times 10^{-8}$).




Figure 16: Mean depth integrated transport streamfunction (10^6 m s^{-1}) for a) the standard calculation; b) a calculation with forcing on the shelf and anticyclonic wind stress curl in the interior. The transport off the shelf is indicated by the black contours (contour interval $= 0.1 \times 10^6 \text{ m s}^{-1}$)




Figure 17: Passive tracer in the model on day 180 at 115 m depth for the case with a wind-forced anticyclonic gyre in the interior.




Figure 18: Mean zonal velocity at 115 m depth for no-slip boundary conditions and free-slip boundary conditions.